


Manipulation of Endophytic Bacteria for Symbiotic Control of *Xylella fastidiosa*, Causal Agent of Citrus Variegated Chlorosis

Paulo T. Lacava
Genetics Department
University of São Paulo/ESALQ
Email: ptlacava@esalq.usp.br
<http://www.esalq.usp.br>




UNIVERSIDADE DE SÃO PAULO
ESCOLA SUPERIOR DE AGRICULTURA "LUIZ DE QUEIROZ"

Citrus Variegated Chlorosis (CVC)



Brazil loses about US\$ 200 million per
year to CVC !!!

Disease Symptoms



Insect Vectors of *X. fastidiosa* (CVC)

Sharpshooter leafhoppers (Homoptera, Cicadellidae, subfamily Cicadellinae)



Acrogonia citrina




Bucephalogonia xanthophis



Dilobopterus costalimai



Oncometopia facialis



Endophytic Microorganisms

Definition

Hallmann et al. (1997): microorganisms that can be isolated from surface disinfected plant tissues and do not induce any visible harm to them

Azevedo et al. (2000): microorganisms inhabit inside plant tissues without neither causing disease to the host nor producing visible external structures.



Host plant

Microorganisms stock center

Epiphytic microorganisms

Endophytic microorganisms

Endophytes colonize ecological niches similar to those colonized by phytopathogens

Application of endophytic microorganisms

Biological control of plant pathogens

- Antimicrobial compounds
- Inducing systemic resistance

5

Interaction between endophytes and plant host: Biotechnological aspects

Wellington Luiz Araújo^{1,2}, Paulo Teixeira Lacava²
Fernando Dini Andreote² and João Lúcio Azevedo^{1,2}

¹Núcleo Integrado de Biotecnologia, Universidade de Mogi das Cruzes, UMC
Avenida Cândido Xavier de Almeida Souza 200, 08780-911, Mogi das Cruzes
SP, Brasil; ²Departamento de Genética, ESALQ/USP, Caixa Postal 9
13418-900, Piracicaba, SP, Brasil

How endophytes could be used as biological control agents of CVC

- ⊕ Select endophytic bacteria to be used in biological control of CVC
- ⊕ Develop a model system to study the interaction between endophytic bacteria and *X. fastidiosa*
- ⊕ CVC control depends on the localization of the endophytic bacteria close to *X. fastidiosa* niche

- **Previous Endophytes Study in Citrus Leaves (Araújo et al. 2001)**

- *Pantoea agglomerans*
- *Bacillus pumilus*

- **Endophytes in Citrus Stems (Araújo et al. 2002)**

- *Methylobacterium spp.*
- *Curtobacterium flaccumfaciens*

Interaction between endophytic bacteria from citrus plants and the phytopathogenic bacteria *Xylella fastidiosa*, causal agent of citrus-variegated chlorosis

P.T. Lacava¹, W.L. Araújo¹, J. Marcon¹, W. Maccheroni Jr¹ and J.L. Azevedo^{1,2}

¹Department of Genetics, Escola Superior de Agricultura 'Luiz de Queiroz', University of São Paulo, Piracicaba, SP, Brazil, and ²Núcleo Integrado de Biotecnologia, University of Mogi das Cruzes, Mogi das Cruzes, SP, Brazil

2003/1097: received 1 December 2003, revised 30 March 2004 and accepted 5 April 2004

ABSTRACT

P.T. LACAVA, W.L. ARAÚJO, J. MARCON, W. MACCHERONI JR AND J.L. AZEVEDO. 2004.


Aims: To isolate endophytic bacteria and *Xylella fastidiosa* and also to evaluate whether the bacterial endophyte community contributes to citrus-variegated chlorosis (CVC) status in sweet orange (*Citrus sinensis* [L.] Osbeck cv. Pera).

Methods and Results: The presence of *Xylella fastidiosa* and the population diversity of culturable endophytic bacteria in the leaves and branches of healthy, CVC-asymptomatic and CVC-symptomatic sweet orange plants and in tangerine (*Citrus reticulata* cv. Blanco) plants were assessed, and the *in vitro* interaction between endophytic bacteria and *X. fastidiosa* was investigated. There were significant differences in endophyte incidence between leaves and branches, and among healthy, CVC-asymptomatic and CVC-symptomatic plants. Bacteria identified as belonging to the genus *Methylobacterium* were isolated only from branches, mainly from those sampled from healthy and diseased plants, from which were also isolated *X. fastidiosa*.

Conclusions: The *in vitro* interaction experiments indicated that the growth of *X. fastidiosa* was stimulated by endophytic *Methylobacterium extorquens* and inhibited by endophytic *Curtobacterium flaccumfaciens*.

Significance and Impact of the Study: This work provides the first evidence of an interaction between citrus endophytic bacteria and *X. fastidiosa* and suggests a promising approach that can be used to better understand CVC disease.

Isolation of culturable endophytic bacteria



Incidence of the main endophytes isolated from branches of sweet orange and tangerine plants

CONCLUSION (Lacava et al , 2004):



- *X. fastidiosa* may colonize CVC-asymptomatic plants without inducing symptoms of CVC;
- *Methylobacterium* spp. were only isolated from branches and mainly from CVC-symptomatic;
- *C. flaccumfaciens* inhibited *in vitro* *X. fastidiosa*. The CVC symptoms could be reduced by a high population of this endophytic bacterium.

Symbiotic Control


- ❖ Relies on the use of symbionts (transformed or not) to reduce vector competence;
- ❖ Pioneered by Frank Richards at Yale University;
- ❖ Trial to suppress Chagas disease through paratransgenesis;
- ❖ Paratransgenesis:
 - * Genetic modification of symbiotic bacteria to make the host less competent in transmitting a pathogen.

Paratransgenesis: The Chagas Disease Case

1- Identify and isolate symbiont



2. Transform symbiont.



3. Reintroduce GM symbiont



Source: slide from J.L. Ramirez

Symbiotic Control

Requirements for a successful paratransgenic approach:

- 1- Symbiont selection
 - * Commensal or mutualistic relationship.
 - * Contact symbiont-pathogen;
- 2- Symbiont culturable and amenable for GM;
- 3- Uncompromised GM symbiont fitness;
- 4- Methods for reintroduction and spread of the GM symbiont.

Beard et al, 2001

CURRENT MICROBIOLOGY Vol. 48 (2004), pp. 327–331
DOI: 10.1007/s00284-003-4178-2

**Current
Microbiology**
An International Journal
© Springer-Verlag New York LLC 2004

Delivery of a Genetically Marked *Alcaligenes* sp. to the Glassy-Winged Sharpshooter for Use in a Paratransgenic Control Strategy

Blake Bextine,¹ Carol Lauzon,² Sarah Potter,² David Lampe,³ Thomas A. Miller¹

¹University of California, Department of Entomology, Riverside, CA 92521-0314, USA

²California State University, 25800 Carlos Bee Blvd., Hayward, CA 94542-3090, USA

³Duquesne University, Biology Department, Pittsburgh, PA 15219, USA

Received: 23 June 2003 / Accepted: 15 August 2003

CURRENT MICROBIOLOGY Vol. 50 (2005), pp. 1–7
DOI: 10.1007/s00284-004-4390-8

**Current
Microbiology**
An International Journal
© Springer Science+Business Media, Inc. 2005

Establishment of a Genetically Marked Insect-Derived Symbiont in Multiple Host Plants


Blake Bextine,¹ David Lampe,² Carol Lauzon,³ Brian Jackson,¹ Thomas A. Miller¹

¹Department of Entomology, University of California, Riverside, CA 92507, USA

²Department of Biological Sciences, Duquesne University, 600 Forbes Avenue, Pittsburgh, PA 15282, USA

³Department of Biological Sciences, California State University, 25800 Carlos Bee Boulevard, Hayward, CA 94542-3090, USA

Received: 3 June 2004 / Accepted: 28 June 2004



The Endophyte *Curtobacterium flaccumfaciens* Reduces Symptoms Caused by *Xylella fastidiosa* in *Catharanthus roseus*

Paulo Teixeira Lacava^{1*}, Wenbin Li², Welington Luiz Araújo¹, João Lúcio Azevedo¹,
and John Stephen Hartung³


¹Department of Genetics, Escola Superior de Agricultura 'Luiz de Queiroz', University of São Paulo, 13400-970 Piracicaba, SP, Brazil

²U.S. Department of Agriculture, Animal and Plant Health Inspection Service Building 580, MD 20705, USA

³U.S. Department of Agriculture-Agricultural Research Service (USDA-ARS), Fruit Laboratory, MD 20705, USA


(Received April 21, 2007 / Accepted September 20, 2007)

RESULTS:



Interaction between *C. flaccunfaciens* (endophytic bacteria) and *X. fastidiosa* inside of host plant (*C. roseus*). Different letters on bars show statistic difference by Tukey's test at 5% of significance

RESULTS:



Interaction between *C. flaccunfaciens* (endophytic bacteria) and *X. fastidiosa* inside of host plant (*C. roseus*). Different letters on bars show statistic difference by Tukey's test at 5% of significance

RESULTS:



Figure 6. Disease symptoms of induced in *Catharanthus roseus* plants 2 months after inoculation with *X. fastidiosa* (right). A symptom free plant doubly-inoculated *X. fastidiosa* and *C. flaccumfaciens* is also shown (left).

RESULTS:



Figure 7. Leaf stunting and chlorosis induced in *Catharanthus roseus* leaves 2 months after inoculation with *Xylella fastidiosa* (left). Symptom free leaves from a plant doubly-inoculated with *X. fastidiosa* and *C. flaccumfaciens* (right).

CONCLUSIONS:

- The results reinforce the idea suggested by Araújo et al. 2002 and Lacava et al. 2004 that this endophytic bacterium, that colonizes a niche similar to that of *X. fastidiosa*, could contribute to the reduction of the symptoms of CVC in the field.

1 **Manipulation of Endophytic Bacteria for Symbiotic Control of Citrus Variegated**

2 **Chlorosis**

3

4 Cláudia Santos Gai^a, Paulo Teixeira Lacava^{a*}, Fernando Dini Andreote^a, Maria Carolina

5 Quecine^a, Marie-Christine Auriac^b, João Roberto Spotti Lopes^a, Wellington Luiz Araújo^{a,c},

6 João Lúcio Azevedo^{a,c}

7

8 ^aEscola Superior de Agricultura "Luiz de Queiroz" – Universidade de São Paulo,

9 Piracicaba, SP, Brazil


10 ^bLaboratoire des Interaction Plantes Microorganismes – INRA/CNRS, Castanet-Tolosan,

11 France

12 ^cNúcleo Integrado de Biotecnologia – Universidade de Mogi das Cruzes, Mogi das Cruzes


13 SP, Brazil

RESULTS:




Plasmidial stability of pCM88 on *Methylobacterium mesophilicum*. The percent was obtained collecting randomly samples after 24, 48, 72 and 120 hours of culture cells of SRGFP strain growing without antibiotic tetracycline.

RESULTS:




Transmission assay. a) Membrane system used for insect feeding, adapted from Purcell and Finlay, 1979. b) Plants with traps containing insects that fed on SRGFP. Arrows indicating traps positioned on plant leaves. c) Detail of trap system.

RESULTS:




Most frequently group of bacteria isolated from *Bucephalogonia xanthophis*.

RESULTS:



Fluorescent microscopy evidencing the ecological niche occupied by endophyte *Methylobacterium mesophilicum* in *Catharanthus roseus* plants 45 days after inoculation.
A) Xylem vessels of a control plant, Scale bar = 10µm; B) Colonized xylem vessel, Scale bar = 5µm; C) Xylem vessels longitudinal cut, Scale bar = 10µm.

RESULTS:



Insects were submitted to isolation form confirmation of the presence of SRGFP colonizing their heads. A) Plate with fluorescent bacteria isolated from insect head after feeding on membrane system. B) Isolation after plant feeding during 24 hours.

Suggestions and Conclusions:

- This is the first report of genetic manipulation of endophytic *M. mesophilicum* isolated from citrus plants;
- *In vitro* the pCM88 plasmid was stably maintained in SRGFP strain for at least 20 generations of cells growing in medium without antibiotic;
- SRGFP strain was stable after the bacteria colonized of plant and insect tissues, therefore representing an important tool to assess both the establishment of the inoculated strain in host plant and the transmission by vector;
- As showed in this study, the transgenic endophytic *M. mesophilicum* has most of the prerequisites for a successful strategy using paratransgenesis (symbiotic control);
- *M. mesophilicum* that colonize citrus plants (Araújo et al. 2002; Lacava et al. 2004) and *B. xanthophilis* is amenable to isolation, culture and transformation with foreign genes.

RESEARCH PAPER

Analysis of the bacterial community in glassy-winged sharpshooter heads

Paulo Teixeira LACAVA¹, Jennifer PARKER², Fernando Dini ANDREOTE¹, Francisco DINI-ANDREOTE¹, José Luiz RAMIREZ² and Thomas A. MILLER²

¹ Department of Genetics, Escola Superior de Agricultura "Luiz de Queiroz", University of São Paulo, Piracicaba, São Paulo, Brazil

² Department of Entomology, University of California Riverside, Riverside, California, USA

- Candidate endophytes for use in SC must occupy the xylem of host plants and attach to the pre-cibarium and cibarium of sharpshooter insects in order to have access to the pathogen;
- The use of isolation and DGGE techniques revealed several genera of bacteria as colonizers of GWSS heads. As identified by 16S sequencing, these included, *Methylobacterium extorquens* and *Curtobacterium flaccumfaciens*;
- Species such as *Curtobacterium flaccumfaciens* and *Methylobacterium* spp., found in part of the bacterial community in GWSS, could be investigated as potential candidates for use in an SC-based strategy to control the spread of PD.

Acknowledgements

- Prof. Dr. João Lúcio de Azevedo, University of São Paulo/ESALQ, SP, Brazil
- Prof. Dr. Welington L. de Araújo, University of São Paulo/ESALQ, SP, Brazil
- Dr. Jonh S. Hartung, USDA-BARC, Beltsville, MD, USA
- Prof. Dr. Thomas A. Miller, University of California Riverside, Riverside, CA, USA
- Prof. João Roberto Spotti Lopes, University of São Paulo/ESALQ, SP, Brazil
- Dr. Cláudia Santos Gai, INRA/CNRS, Castante - Tolosan, France
- The Brazilian agency FAPESP

